
Climate Variability, Hydrology, and Flooding



Overview of NASA Hydrological Data Precipitation, Snow Cover



Objective

To provide an overview of precipitation products (rain and snow) from NASA satellites and models. Precipitation is major component of freshwater, responsible for extreme weather and flooding and associated with regional climate variability

Outline

- Precipitation data from NASA Satellites and Models
- Rainfall Data and Access

The Tropical Rainfall Measurement Mission (TRMM)
The Global Precipitation Measurement (GPM) Mission
TRMM Applications

Snow Data and Access

The Aqua/Terra Moderate Resolution Imaging Spectroradiometer (MODIS)

Precipitation data from NASA Satellites and Models

NASA Remote Sensing Data for Rain and Snow

Satellite	Sensors	Quantities
TRMM	Precipitation Radar (PR) TRMM Microwave Imager (TMI) Visible Infrared Scanner (VIRS)	Rain Rate, Vertical Rain Rate Profile, Accumulated Rain
GPM	Dual Frequency Precipitation Radar (DPR) GPM Microwave Imager (GMI)	Rain Rate, Vertical Rain Rate Profile, Accumulated Rain
Terra and Aqua	MODerate Resolution Imaging Spectroradiometer (MODIS)	Snow Cover, Vegetation Index, Leaf Area Index, Land Cover

Rain Rate has unit of mm/hour Accumulated Rain has unit of mm (over a day or a month) Snow Cover is the fractional area covered by snow

NASA Models for Precipitation

	Models	Quantities
These models assimilate satellite observations and are useful for weather, climate, and hydrology	MERRA	3-dimensional Winds, Temperature, Humidity, Clouds, Rain Rate, Snow Mass, Snow Cover, Snow Depth, Surface Snowfall Rate, Evapotranspiration
research This training will focus on higher spatial resolution satellite observations for rain and snow data	GLDAS	Evapotraspiration, Multi-layer Soil Moisture, Rainfall, Snowfall Rate, Snow Melt, Snow-Water Equivalent, Surface and Sub-surface Runoff

Snow Rate has unit of mm/hour Snow Water Equivalent has units of kg/m² or m

Overview of Tropical Rainfall Measurement Mission (TRMM) Data and Access

Important Note:

TRMM mission was terminated in April 2015 but near-real time TRMM-calibrated rainfall from other satellites are available until GPM data become available in near-real time

TRMM data from 1997-2014 are widely used for weather, climate, and hydrology applications

Review of TRMM Tropical Rainfall Measuring Mission

http://trmm.gsfc.nasa.gov


- The first satellite mission dedicated to measuring tropical and subtropical rainfall - Launched on 27 November 1997
- First satellite to carry a microwave Precipitation Radar
- Predecessor to Global Precipitation Measurement (GPM)



Review of TRMM

http://trmm.gsfc.nasa.gov


- A non-polar, low inclination orbit
 Revisit time ~11-12 hours, but time of the observation changes daily
- There are 16 TRMM orbits a day covering global tropics between 35° S to 35°N latitudes
- Altitude of approximately 350 Km,
 raised to 403 Km after 23 August 2001



TRMM

http://trmm.gsfc.nasa.gov

- Multiple sensors
- One active and two passive rain sensors
 - Precipitation Radar (**PR**)
 TRMM Microwave Imager (**TM**I)
 Visible and Infrared Scanner (**VIRS**)
- Multiple rain products available from individual sensors, at varying spatial resolutions



TRMM Multi-satellite Precipitation Analysis (TMPA)

(Widely used in Environmental Applications)

Combination of TRMM - TMI, PR, VIRS with passive microwave, infrared and visible measurements available from national and international satellites provides rainfall data with --

Temporal Resolution: 3-hourly

Spatial Resolution: 0.25°x0.25°

Spatial Coverage: Global 50°S to 50°N

Available in Near-real time data and also gauge-calibrated research quality version with ~ 3-month latency

This training will focus on TMPA because it is available in near-real time and also for 15+ years – useful for flooding and climate applications

Summary of TRMM Level-2 Precipitation Products

*Surface Rainfall Rate in mm/hour

TRMM data are available from December 1997 to present

Sensor/Product Name	Spatial Resolution and Coverage	Temporal Resolution	Data Format
TMI/2A12	5 km x 5 km Orbital and 16 orbits per day (38°S-38°N)	3-hour, 2-day 15 days	HDF4 and OPenDAP
PR/2A25	5 km x 5 km Orbital and 16 orbits per day (38°S-38°N)	3-hour, 2-day 5 days	
Combined TMI and PR /2B31	5 km x 5 km Orbital and (38°S-38°N)	3-hour, 2-day 5 days	

^{*}In addition to surface rainfall rate in mm//hour, vertical precipitation profiles and latent heating are available in these data products

Summary of TRMM Level-3 Precipitation Products

*Surface Rainfall Rate in mm/hour


TRMM data are available from January 1998 to present

Sensor/Product Name	Spatial Resolution and Coverage	Temporal Resolution	Data Format
TMPA/3B42 RT	0.25°x0.25° (50°S-50°N)	3-hourly (Near Real Time) daily, 10-day and 30-day Averages	HDF4, NetCDF, OPenDAP, ASCII GIF, PNG Images KML for Google Earth
TMPA/3B42 (Gauge Adjusted Research Version)	0.25°x0.25° (50°S-50°N)	3-hourly, daily	
TMPA/3B43	0.25°x0.25° (50°S-50°N)	Monthly	

^{*}In addition to surface rainfall rate in mm//hour, vertical precipitation profiles and latent heating are available in these data products


TRMM TMPA Near-Real Time Images

http://pmm.nasa.gov/TRMM/realtime-3hr-7day-rainfall



TRMM TMPA Access through Giovanni-4


http://giovanni.gsfc.nasa.gov/giovanni/



Daily and Monthly 0.25 degree TMPA (Product names: 3B42 and 3B43) available in NetCDF format, PNG or GeoTIFF Images

TRMM TMPA Access through Mirador

http://mirador.gsfc.nasa.gov/



Mirador is useful for searching data and downloading multiple data files in HDF or OpenDAP Formats

Global Precipitation Measurement Mission (GPM) Designed to extend, enhance, and improve TRMM Precipitation Data

TRMM Data Limitations:

Does not provide measurements beyond 35°S-35°N

The TRMM sampling frequency is 15 hours to 4 days at any point which can introduc substantial uncertainties in rain estimates

TRMM provides rain measurements but not frozen precipitation, also can not detect light rain (<0.5 mm/hr)

GPM was designed to obtain measurements over tropics and higher latitudes, with improved capability to observe light rain and snow


Overview of Global Precipitation Measurement (GPM) Data and Access

Global Precipitation Measurement (GPM)

http://pmm.nasa.gov/GPM


- An international network of satellites with a GPM Core satellite designed to provide global observations of rain and snow
- Initiated by NASA and the JAXA as a successor to TRMM

GPM Core satellite was launched on February 27th, 2014 GPM Constellation Status



GPM Orbits and Spatial Coverage

- In non-polar, low inclination orbit with 16 orbits per day
- GPM observes global regions between 65°
 S to 65°N latitudes



the area covered by three TRMM orbits [yellow] versus orbits of the GPM Core Observatory [blue]

GPM measurements span middle and high latitudes

GPM GMI and DPR Measurements

http://pmm.nasa.gov/GPM

GMI

DPR

Compared to TRMM TMI:

- Higher spatial resolutions
- Improved light rain and snow detection
- Reference for constellation radiometers calibration

Compared to TRMM PR:

- Higher sensitivity to light rain and snow
- Better accuracy of measurements
- Better identification of liquid, ice, mixed-phase precipitation particles
- Reference standard for intercalibration of constellation precipitation measurements

Summary of GPM Level-2 Precipitation Products

*Surface Rainfall Rate in mm/hour

GPM data are available from March 2014 to present

Sensor/Product Name	Spatial Resolution and Coverage	Temporal Resolution	Data Format
DPR Ku-only/ 2A-Ku DPR Ka-only/2A-Ka DPR KU & Ka/ 2A-DPR	5.2 km x125 m Single Orbit and 16 orbits per day (70°S-70°N)	20-120 minutes 24 hours	HDF5 and OPenDAP
GMI/2A-GPROF	4 km x 4 km Orbital and 16 orbits per day (70°S-70°N)	2 – 40 hours	
Combined GMI and DPR/2A-CMB	Orbital (70°S-70°N) 5 km x 5 km, Coincident Ku-Ka-GMI footprints	3 – 40 hours	

^{*}In addition to surface rainfall rate in mm//hour, vertical precipitation profiles and latent heating are available in these data products

Summary of GPM Level-3 Precipitation Products

*Surface Rainfall Rate in mm/hour

GPM data are available from March 2014 to present

Sensor/Product Name	Spatial Resolution and Coverage	Temporal Resolution	Data Format	
IMERG	0.1°x0.1° (90°S-90°N)	30-minutes(Near Real Time) with 4-hour latency, 12-hour latency and 4- months latency	HDF4, NetCDF, OPenDAP, ASCII GIF, PNG Images KML for Google Earth	
3-CMB Combined GMI + DPR rainfall Averages	0.1°x0.1° (70°S-70°N)	Monthly		
3-DPR rainfall Averages	0.25°x0.25° 5.0°x5.0° (67°S-67°N) for Daily (70°S-70°N) for Monthly	Daily and Monthly Daily and Monthly		
3-GPROF GMI rainfall Averages	0.25°x0.25° (90°S-90°N)	Daily and Monthly		

^{*}In addition to surface rainfall rate in mm//hour, vertical precipitation profiles and latent heating are available in these data products

TMPA: TRMM Multi-satellite Precipitation Analysis IMERG: Integrated Multi-satellitE Retrievals for GPM

IMERG is Conceptually similar to TRMM TMPA, combines GPM GMI/DPR data with the GPM constellation satellites to yield improved spatial/temporal precipitation estimates:

IMERG TMPA

Temporal Resolution: 30-minutes 3 hours

Spatial Resolution: 0.1°x0.1° 0.25°x0.25°

Spatial Coverage: Global Global

60°S to 60°N 50°S to 50°N

Constellation Satellites:

2. IMERG Data Sets

Multiple runs accommodate different user requirements for latency and accuracy

- •"Early" 4 hours (flash flooding)
- •"Late" 12 hours (crop forecasting)
- "Final" 3 months (research data)

Time intervals are half-hourly and monthly (Final only)

0.1° global CED grid

- PPS will provide sub-setting by parameter and location
- initial release covers 60°N-S

User-oriented services

- interactive analysis (GIOVANNNI)
- alternate formats (KMZ, KML, TIFF WRF files, ...)
- area averages


	Half-hourly data file (Early, Late, Final)	
1	[multi-sat.] precipitationCal	
2	[multi-sat.] precipitationUncal	
3	[multi-sat. precip] randomError	
4	[PMW] HQprecipitation	
5	[PMW] HQprecipSource [identifier]	
6	[PMW] HQobservationTime	
7	IRprecipitation	
8	IRkalmanFilterWeight	
9	probabilityLiquidPrecipitation [phase]	
	Monthly data file (Final)	
1	[satgauge] precipitation	
2	[satgauge precip] randomError	
3	GaugeRelativeWeighting	
4	probabilityLiquidPrecipitation [phase]	

GPM Data Access Tools

	Tools	Data Products and Formats	Analysis and/or Visualization	Data Download
	Mirador http://mirador.gsfc.nasa.gov Same as for	L1B, L2, and L3 GMI-GPROF IMERG Half-hourly, Monthly Orbital and Gridded Daily, Monthly HDF5, OPenDAP (can be converted to ASCII, Binary, NetCDF)		Batch Download
	Giovanni http:// giovanni.gsfc.nasa.gov/ giovanni/	IMERG Half-hourly, Monthly NetCDF, GeoTIFF, PNG	Visualization: Map, Time Series, Scatter Plot Histogram Analysis: Time-averaged Maps, Time Series, Scatter Plot, Map Correlations, Vertical Profiles, Time- averaged Differences	Download by Select and Click on Data Files
	PPS/STORM https:// storm.pps.eosdis.nasa.gov/ storm	L1B and 1C, L2, L3 GMI, DPR, GMI-DPR Combined Data, Orbital and Gridded Daily, Monthly IMERG Half-hourly, Monthly HDF5, PNG	Map Visualization, Interactive Latitude/Longitude Point Data Value Display	FTP


26

https://storm-pps.gsfc.nasa.gov/storm/




STORM is specifically designed for GPM and TRMM Precipitation data search, selection, download, and visualization

https://storm-pps.gsfc.nasa.gov/storm/



STORM is specifically designed for GPM and TRMM Precipitation data search, selection, download, and visualization

https://storm-pps.gsfc.nasa.gov/storm/




https://storm-pps.gsfc.nasa.gov/storm/



https://storm-pps.gsfc.nasa.gov/storm/

Product Selection, Download, and Visualization by using Tool for High-resolution Observation Review (THOR)



Precipitation on November 15, 2014

TRMM data are used for a variety of applications, these applications will continue, using improved GPM data

PRECIPITATION DATA APPLICATIONS

Societal Benefit Areas of TRMM and GPM Precipitation



Extreme Events and Disasters



Water Resources and Agriculture



Weather, Climate & Land Surface Modeling

- Numerical Weather Prediction
- Land System Modeling Global ClimateModeling




Public Health and Ecology

TRMM Data Are Used in Weather Monitoring

Used by AccuWeather (http://www.accuweather.com) to monitor storms and heavy rainfall

Monster Australia Cyclone Ita, April 9 & 10, 2014

- GPM data will be used by tropical cyclone forecasting centers worldwide to detect the location and intensity of tropical cyclones.
- GPM's orbit (unlike TRMM's) will enable observation of tropical cyclones as they progress from tropical to mid-latitude systems



TRMM PR and TMI showing heavy rainfall within the storm

TRMM Data Are Used to Provide Early Warning of Extreme Rainfall and Flooding For Developing Countries

GPM-IMERG will be used for extreme rainfall detection by ITHACA)

Used by Information Technology for Humanitarian Assistance, Cooperation, and Action (ITHACA) www.ithacaweb.org

Extreme Rainfall Detection System –
 Version 2 (ERDS2) uses near-real time
 3-hourly TMPA

(http://www.ithacaweb.org/projects/erds/)

ERDS2 is a strategic tool, providing immediate information about potential flood events, used by the UN World Food Programme (WFP) Emergency Preparedness Unit

ITHACA provides trainings to government staff in developing countries on how to use ERDS2 and remote sensing data for flood hazard assessment

http://www.ithacaweb.org/news/



ITHACA delivers technical training to Malawi Gov't staff

Fri 16 Jan 2015

Within the World Bank financed project MASDAP, ITHACA held a technical training session on the use of satellite data for vulnerability assessment in Blantyre from December 15 to 19, 2014.

Read more.

ITHACA hosts 3 Ethiopian interns for technical training on GIS and GPS systems


Mon 20 Oct 2014

On October 20 ITHACA started a 3-week specialization module on GIS and GPS systems for 3 Ethiopian trainees in the framework of the WATSAM project, coordinated by Hydroaid - Water for Development Management Institute. Read more.

Satellite-based Rapid Mapping training in Lilongwe₃₅ (September 29 – October 1, 2014)

Wed 15 Oct 2014

TRMM Data Are Used for Flooding Estimates



TRMM Data Are Used by Global Disaster Alert and Coordination System (GDACs)

GDACS, managed by the United Nations and European Commission, provides disaster alerts and river watch that is used by many governments and about 14,000 disaster response and non-governmental organizations for their national disaster response plans. (http://www.gdacs.org)

One of the data sources used by GDACS is river run off derived from TRMM-TMI data by Dartmouth Flood Observatory


(http://www.dartmouth.edu)

Higher resolution and extended spatial coverage by GPM-GMI will provide improved river-run off estimates



Flooding alert in Botswana on 15th December 2014

TRMM Data Are Used in Agricultural Forecasting



http://www.fews.net

Famine Early Warning System (FEWS) relies on TRMM and other satellite estimates for anticipating poor growing seasons. GPM will improve these estimates.

FEWS NET Data Portal

http://earlywarning.usgs.gov/fews






TRMM Data Are Used in Disease Tracking

TRMM data has been used to estimate and trace the source areas of vector and river-borne diseases around the world. GPM will enable higher resolution evaluation of these disease source areas.


Observed Plague Cases in Uganda



Cases are associated with wetter, cooler regions

Monaghan et al. 2012; MacMillan et al., 2012

Schistosomiasis (snail-spread) in Ethiopiamiasis



Courtesy of Bitew and

Studies have found a relationship between TRMM rain and the onset of this disease in local populations due to contact with snails in irrigation channels


Snow Data and Access

Review of Terra and Aqua

http://trmm.gsfc.nasa.gov

Terra


- Polar, Sun-Synchronous Orbit, Global Coverage
- Twice-daily Observations 10:30 AM/PM Descending Orbits
- From 12/1999 Present



Local noon (ascending) AIRS CO at 500 mb on 2010,08.01.

Aqua

- Polar, Sun-Synchronous Orbit, Global Coverage
- Twice-daily Observations 1:30 AM/PM Descending Orbits
- From 5/2002 Present



Terra and Aqua MODerate Resolution ImagingSpectroradiometer (MODIS)

http://modis.gsfc.nasa.gov/

- A key instrument aboard Terra ad Aqua providing 4-times per day observations (1:30 and 10:30 AM/ PM) from the two satellites
- There are 36 spectral bands designed to observe atmosphere, land, and ocean
- Multiple data products and tools available

More information about obtaining MODIS data can be found from the information sites listed below. The URL's for these sources of data are as follows:

MODIS Level 1 data, geolocation, cloud mask, and Atmosphere products:

http://ladsweb.nascom.nasa.gov/

MODIS land products:

https://lpdaac.usgs.gov/

MODIS cryosphere products:

http://nsidc.org/daac/modis/index.html


MODIS ocean color and sea surface temperature products:

http://oceancolor.gsfc.nasa.gov/



Terra and Aqua MODerate Resolution Imaging Spectroradiometer (MODIS)

- 36 spectral bands ranging from 0.41 to 14.385 microns
- Many applications, including clouds, snow/ ice, vegetation, aerosol
- Available in various resolution (depends on product)





http://modis-snow-ice.gsfc.nasa.gov/

There are Two MODIS-based Snow Data Products

- Standard MODIS Product: Fractional Snow Cover
- MODSCAG (MODIS Snow Covered Area and Grainsize) Product : Fractional Snow Cover, Grain Size, Snow Water Equivalent

Snow Dominated Regions




Table 2. MODIS snow data products.

From Dorothy Hall NASA-GSFC

Long Name	Earth Science Data Type (ESDT)	Spatial Resolution
MODIS/Terra Snow Cover 5-Min L2 Swath 500m	MOD10_L2	500-m resolution, swath of MODIS data
MODIS/Terra Snow Cover Daily L3 Global 500m SIN Grid (includes daily snow albedo)	MOD10A1	500-m resolution, projected, gridded tile data
MODIS/Terra Snow Cover 8-Day L3 Global 500m SIN Grid	MOD10A2	500-m resolution, projected, gridded tile data
MODIS/Terra Snow Cover Daily L3 Global 0.05Deg CMG	MOD10C1	0.05° resolution, lat/lon climate modeling grid
MODIS/Terra Snow Cover 8-Day L3 Global 0.05Deg CMG	MOD10C2	0.05° resolution, lat/lon climate modeling grid
MODIS/Terra Snow Cover Daily L3 Global 0.25Deg CMG	Not yet a standard product	0.25° resolution, lat/lon climate modeling grid
MODIS/Terra Snow Cover Monthly L3 Global 0.05Deg CMG	MOD10CM	0.05° resolution, lat/lon climate modeling grid
MODIS/Aqua Snow Cover 5-Min L2 Swath 500m	MYD10_L2	500-m resolution, swath of MODIS data
MODIS/Aqua Snow Cover Daily L3 Global 500m SIN Grid (includes daily snow albedo)	MYD10A1	500-m resolution, projected, gridded tile data
MODIS/Aqua Snow Cover 8-Day L3 Global 500m SIN Grid	MYD10A2	500-m resolution, projected, gridded tile data
MODIS/Aqua Snow Cover Daily L3 Global 0.05Deg CMG	MYD10C1	0.05° resolution, lat/lon climate modeling grid
MODIS/Aqua Snow Cover 8-Day L3 Global 0.05Deg CMG	MYD10C2	0.05° resolution, lat/lon climate modeling grid
MODIS/Aqua Snow Cover Monthly L3 Global 0.05Deg CMG	MYD10CM	0.05° resolution, lat/lon climate modeling grid

MODIS Standard Snow Product Data Access and Visualization

MODIS Snow Products are available from the National Snow and Ice Data Center http://nsidc.org/

MODIS monthly snow cover can be visualized on Google Earth maps from http://nsidc.org/data/virtual_globes/index.html

Data Start Date: 2000-02-24

Daily, 8-Day, Monthly


Coverage: Global

Multiple Spatial Resolutions




Selected Aqua-MODIS Snow Product from Reverb/ECHO

http://reverb.echo.nasa.gov/reverb



Near-real Time Standard MODIS Products

http://lance-modis.eosdis.nasa.gov/



Near-real Time Standard MODIS Products

http://lance-modis.eosdis.nasa.gov/

Level-2 Swath data (500) m and 5 Km resolution Snow Cover available from Terra and Aqua MODIS

Terra


			Fraction Browse					
L2 Snow Cover, 5-Min Swath 500m	MOD10_L2	0.26	L2 Snow Cover Browse	N/A	07	0:46	1:32 (8)	3:14
L2 Coarse Snow Cover, 5-Min Swath 5km	MOD10L2C	0.17				0:46	1:32 (8)	3:14
L2 Coo los Extent							1:20	

Aqua

			Browse					
L2 Snow Cover, 5-Min Swath 500m	MYD10_L2	0.26	L2 Snow Cover Browse	N/A	07	1:00	1:47 (25)	3:30
L2 Coarse Snow Cover, 5-Min Swath 5km	MYD10L2C	0.17				1:00	1:47 (25)	3:30


MODSCAG Snow Products

From: Thomas H. Painter and Chris Mattmann (NASA JPL)




MODSCAG

http://snow.jpl.nasa.gov/portal/browse/dataset/urn:snow:MODSCAG



Limitation of MODIS Data: No Snow Mapping Under Clouds

http://snow.jpl.nasa.gov/portal/browse/dataset/urn:snow:MODSCAG



Water Resources



Integration of Precision NASA Snow Products with the Operations of the Colorado Basin River Forecast Center (CBRFC) to Improve Decision Making Under Drought Conditions

Principle Investigator: Thomas Painter, Jet Propulsion Laboratory

Abstract

The Colorado Basin River Forecast Center (CBRFC) is responsible for the entire Colorado Basin (CRB) and the eastern Great Basin (GB). From a water management perspective, the commitment of water to various users most often occurs in the spring, and is almost entirely based on estimates of the western USA snowpack. Improving seasonal drought predictions requires use of models that provide physically realistic simulations of fundamental hydrologic processes. Among these, for the western USA, representation of snow is perhaps most critical.

As drought frequency increases in the CRB and GB, it is critical that the CBRFC and the dependent water managers have more comprehensive real-time knowledge of the snow cover and its properties for more precise runoff forecasting and stakeholder decision support. The primary objective of this proposal is to integrate real-time high precision MODIS Snow Covered Area and Grain size (MODSCAG) fractional snow covered area (SCA) into CBRFC modeling and analysis systems and into stakeholder oriented data products, drastically reducing SCA uncertainties that have hampered forecasting operations for decades. A secondary objective is to ingest and study MODIS Dust Radiative Forcing in Snow (MODDRFS) radiative forcing imagery, to better understand its value as a input to modeling and forecasting approaches.

This collaboration directly addresses drought prediction, assessment, adaptation, and mitigation in support of energy security/efficiency; natural resource conservation; and household, municipal, industrial, and in-stream demands for water. It will also improve access and availability of actionable water monitoring, hence drought information. he Snow Cover and Dust Forcing products will be generated and distributed in near real-time by the JPL Snow Server for access by CBRFC. CBRFC will offer a direct connection to stakeholders (End Users) and together with other linked NWS operational centers provides an institutional home to maintain the advances of this effort beyond the project's end.


MODSCAG, Snow and Dust Radiative Forcing Information, along with CBRFC Modeling Analysis is used in Decision Making for River Basin management

top 53

Next:

Hands-on Activity to Access, Visualize, Download Monthly TMPA Data using Giovanni-4


TRMM TMI and PR Measurements



Strength: High pixel resolution, Accurate measurements **Limitation**: **No global, diurnal coverage on daily basis**

GPM GMI and DPR Measurements

http://pmm.nasa.gov/GPM




GMI Frequencies: 10.6,18.7,23.8,36.5,89,166 & 183 GHz


Swath width 885 km


Resolution: 19.4km x 32.2km (10 GHz)

to 4.4km x 7.3km (183 GHz)

Higher spatial resolutions than TMI High frequencies help measure snow







Ku 13.6 GHZ, Swath Width 245 km, Resolution 5.2 km